

| Specie | Immediate Epigeal | Delayed Epigeal | Delayed Hypogeal | Autumnal/ Winter Hypogeal | Comments |
|---|-------------------|-----------------|------------------|---------------------------|--|
| <i>L. alexandrae</i> syn. <i>L. ukeyuri</i> | | | * | | Usually exhibits autumnal germination. |
| <i>L. amabile</i> | * | | | | |
| <i>L. amoenum</i> | * | | | | |
| <i>L. albanicum</i> | | | | | See <i>L. pyrenaicum</i> ssp. <i>carniolicum</i> var. <i>albanicum</i> |
| <i>L. apertum</i> | * | | | | Synonym <i>Nomocharis aperta</i> |
| <i>L. arboricola</i> | * | | | | |
| <i>L. armenum</i> | | | * | | |
| <i>L. auratum</i> | | | * | | |
| <i>L. bakerianum</i> | * | | | | |
| <i>L. bolanderi</i> | | | | * | |
| <i>L. bosniacum</i> | | | | | See <i>L. pyrenaicum</i> ssp. <i>carniolicum</i> var. <i>bosniacum</i> |
| <i>L. brownii</i> | * | | | | |
| <i>L. bulbiferum</i> | | | * | | |
| <i>L. callosum</i> | * | | | | |
| <i>L. columbianum</i> | | | | * | |
| <i>L. canadense</i> | | | * | | |
| <i>L. candidum</i> | * | | | | |
| <i>L. carniolicum</i> | | | | | See <i>L. pyrenaicum</i> ssp. <i>carniolicum</i> |
| <i>L. catesbaei</i> | * | | | | |
| <i>L. cernuum</i> | * | | | | |
| <i>L. chalcedonicum</i> | | * | | | |
| <i>L. ciliatum</i> | | | * | | |
| <i>L. concolor</i> | * | | | | |
| <i>L. dauricum</i> | * | | | | |
| <i>L. davidii</i> | * | | | | |
| <i>L. distichum</i> | | | * | | |
| <i>L. duchartrei</i> | * | | | | |
| <i>L. eupetes</i> | * | | | | |
| <i>L. fargesii</i> | * | * | | | A few germinate immediately more germinate after a cold delay period. |
| <i>L. formosanum</i> | * | | | | |
| <i>L. gloriosoides</i> (Chinese form) | | | | | See <i>L. speciosum</i> var. <i>gloriosoides</i> (Chinese Form) |
| <i>L. gloriosoides</i> (Taiwanese form) | | | | | See <i>L. speciosum</i> var. <i>gloriosoides</i> (Taiwanese Form) |
| <i>L. grayi</i> | | | * | | |
| <i>L. hansonii</i> | | | * | | |

| | | | | | |
|--|---|--|---|---|---|
| <i>L. henrici</i> | * | | | | |
| <i>L. henryi</i> | * | | | | |
| <i>L. humboldtii</i> | | | | * | Some southern forms exhibit immediate autumnal germination. |
| <i>L. iridollae</i> | | | * | | |
| <i>L. jankae</i> | | | | | See <i>L. pyrenaicum</i> ssp. <i>carniolicum</i> var. <i>jankae</i> |
| <i>L. japonicum</i> | | | * | | |
| <i>L. jinfushanense</i> | * | | | | |
| <i>L. kelleyanum</i> | | | | * | |
| <i>L. kelloggii</i> | | | | * | |
| <i>L. kesselringianum</i> | | | * | | |
| <i>L. lancifolium</i> | * | | | | |
| <i>L. lankongense</i> | * | | | | |
| <i>L. leichtlinii</i> | * | | | | |
| <i>L. ledebourii</i> | | | * | | |
| <i>L. leucanthum</i> | * | | | | |
| <i>L. lijiangense</i> syn. <i>L. tenii</i> | * | | | | |
| <i>L. longiflorum</i> | * | | | | |
| <i>L. mackliniae</i> | * | | | | |
| <i>L. maculatum</i> | * | | | | |
| <i>L. majoense</i> | * | | | | |
| <i>L. maritimum</i> | | | | * | Often appear above ground in late winter. |
| <i>L. martagon</i> | | | * | | |
| <i>L. matangense</i> | * | | | | |
| <i>L. medogense</i> | * | | | | |
| <i>L. medeoloides</i> | | | * | | |
| <i>L. meleagrinum</i> | * | | | | Synonym <i>Nomocharis meleagrina</i> |
| <i>L. michauxii</i> | | | * | | Texan populations appear above ground in autumn. |
| <i>L. michiganense</i> | | | * | | |
| <i>L. monadelphum</i> | | | * | | |
| <i>L. nanum</i> | * | | | | |
| <i>L. neilgherrense</i> | * | | | | |
| <i>L. nepalense</i> | * | | | | |
| <i>L. nobillissimum</i> | | | * | | Usually exhibits autumnal germination. |
| <i>L. occidentale</i> | | | | * | |
| <i>L. oxypetalum</i> | * | | | | |
| <i>L. papilliferum</i> | * | | | | |
| <i>L. paradoxum</i> | * | | | | |

| | | | | | |
|--|---|---|---|---|---|
| <i>L. pardalinum</i> | | | | * | |
| <i>L. pardalinum</i> ssp. <i>pitkinense</i> | | | | * | |
| <i>L. pardalinum</i> ssp. <i>shastense</i> | | | | * | |
| <i>L. pardalinum</i> ssp. <i>vollmeri</i> | | | | * | |
| <i>L. pardalinum</i> ssp. <i>wigginsii</i> | | | | * | |
| <i>L. pardanthinum</i> | * | | | | Synonym <i>Nomocharis pardanthina</i> |
| <i>L. parryi</i> | | | | * | May exhibit immediate hypogeal germination. |
| <i>L. parvum</i> | | | | * | |
| <i>L. philadelphicum</i> | * | | | | |
| <i>L. philippinense</i> | * | | | | |
| <i>L. pitkinense</i> | | | | * | See <i>L. pardalinum</i> ssp. <i>pitkinense</i> |
| <i>L. polyphyllum</i> | | | * | | |
| <i>L. pomponium</i> | | * | | | Often appear above ground in late winter. |
| <i>L. ponticum</i> | | * | | | |
| <i>L. primulinum</i> | * | | | | |
| <i>L. pumilum</i> | * | | | | |
| <i>L. pyrenaicum</i> | | * | | | |
| <i>L. pyrenaicum</i> subsp. <i>carniolicum</i> | | * | | | |
| <i>L. pyrenaicum</i> subsp. <i>carniolicum</i> var. <i>albanicum</i> | | * | | | |
| <i>L. pyrenaicum</i> subsp. <i>carniolicum</i> var. <i>bosniacum</i> | | * | | | |
| <i>L. pyrenaicum</i> subsp. <i>carniolicum</i> var. <i>jankae</i> | | * | | | |
| <i>L. pyrophyllum</i> | | | * | | |
| <i>L. regale</i> | * | | | | |
| <i>L. rhodopaeum</i> | | * | | | |
| <i>L. rosthornii</i> | * | | | | |
| <i>L. rubellum</i> | | | * | | |
| <i>L. rubescens</i> | | | | * | |
| <i>L. sargentiae</i> | * | | | | |
| <i>L. sealyi</i> | * | | | | Synonym <i>Nomocharis farreri</i> |
| <i>L. shastense</i> | | | | | See <i>L. pardalinum</i> ssp. <i>shastense</i> |
| <i>L. sherriffiae</i> | * | | | | |
| <i>L. soulei</i> | * | | | | |
| <i>L. speciosum</i> | | | * | | |
| <i>L. speciosum</i> var. <i>gloriosoides</i> (Chinese Form) | | | * | | |
| <i>L. speciosum</i> var. <i>gloriosoides</i> (Taiwanese Form) | * | | | | |

| | | | | | |
|---|---|--|---|---|--|
| <i>L. stewartianum</i> | * | | | | |
| <i>L. sulphureum</i> | * | | | | |
| <i>L. superbum</i> | | | * | | |
| <i>L. szovitsianum</i> | | | * | | |
| <i>L. taliense</i> | * | | | | |
| <i>L. tsingtauense</i> | | | * | | |
| <i>L. ukeyuri</i> syn. <i>L. alexandrae</i> | | | * | | Usually exhibits autumnal germination. |
| <i>L. vollmeri</i> | | | | * | See <i>L. pardalinum</i> ssp. <i>vollmeri</i> |
| <i>L. wallichianum</i> | * | | | | |
| <i>L. washingtonianum</i> | | | | * | |
| <i>L. wardii</i> | * | | | | |
| <i>L. wenshanense</i> | | | * | | Usually exhibits autumnal germination. |
| <i>L. wigginsii</i> | | | | * | See <i>L. pardalinum</i> ssp. <i>wigginsii</i> |
| <i>L. wilsonii</i> syn. <i>L. maculatum</i> | * | | | | |
| <i>L. yapingense</i> | * | | | | |